


Chapter 7 TREATMENT OF SPECIFIC HEMORRHAGES

Johnny Mahlangu, Gerard Dolan, Alison Dougall, Nicholas J. Goddard, Enrique D. Preza Hernández, Margaret V. Ragni, Bradley Rayner, Jerzy Windyga, Glenn F. Pierce, Alok Srivastava

Treatment of Specific Hemorrhages: WHAT YOU NEED TO KNOW



1.Intracranial hemorrhage

Symptoms:

- Head injury
- Headache
- Sleepiness
- Severe back pain

Treatment:

- Immediate treatment with CFCs (before investigations begin)
- Immediate hospitalization and medical evaluation by a neurologist

Gastrointestinal and abdominal hemorrhage

Symptoms:

- Vomiting blood
- Rectal passage of blood
- Melena

Treatment:

- Immediate CFCs (before investigations begin)
- Immediate hospitalization and medical evaluation
- Antifibrinolytics
- Treat underlying condition

2.Renal hemorrhage (bleeding in kidneys)

Symptoms:

- Abdominal pain
- Abdominal swelling
- Flank and back pain
- Urination of blood

Treatment:

- Immediate treatment with CFCs
- Adequate hydration
- Bed rest

4. Joint hemorrhage

Symptoms:

- Tingling sensation and tightness within the joint
- Swelling, skin warm to the touch
- Increase of pain
- Loss of range of motion

Treatment:

- Immediate CFCs
- Analgesics for pain
- RICE (rest, ice, compression, elevation), PRICE (protection, rest, ice, compression, elevation), or POLICE (protection, optimal loading, ice, compression, elevation)
- Physical therapy and rehabilitation



CFC = clotting factor concentrate replacement therapy

Chapter 7 TREATMENT OF SPECIFIC HEMORRHAGES



1.Ophthalmic hemorrhage

(bleeding in the eye)

Cause:

• Possible trauma or infection

Treatment:

- Immediate treatment with CFCs
- Monitor every 6-8h until bleed has resolved

2. Epistaxis (bleeding from the nose)

Cause:

- Irritation
- Trauma

Treatment:

- Elevate head
- Compression
- Gauze soaked in antifibrinolytics
- Avoid nasal packing

4.Throat and neck

• CFCs if warranted



3.Oral hemorrhage

Cause:

- Dental extraction
- Gum disease
- Trauma

Treatment:

- Apply direct pressure 15-30 min / sutures and adrenaline if needed
- Immediate treatment with CFCs
- Dentist consultation
- No intense sports, soft diet x3-5 days
- No smoking for 24 hrs
- Soft splint if necessary

hemorrhages Cause:

- Local pathology
- Trauma
- Severe coughing

Treatment:

- Immediate treatment with CFCs
- Immediate hospitalization and medical evaluation by an otolaryngologist



